



Indian Veterinary Magazine

a monthly Magazine

of the people, by the people, for the people



Vol 2 Issue 2 Feb 2026, 110-115

ISSN:3108-1398

Low-Cost Feeding Strategies for Dairy Farmers in India

^aPramod Kumar, ^bNeha Sahu, ^aSuchit Kumar, ^bVinay V V

Animal Nutrition Division, ICAR-National Dairy Research Institute, Karnal, Haryana
Assistant Professor, Livestock Farm Complex, Bihar Veterinary College, BASU, Patna

Introduction

Dairy farming plays an important role in the livelihood of millions of farmers in India, providing regular income, employment, and nutritional security. However, one of the major challenges faced by dairy farmers is the high cost of feeding dairy animals. Feed alone accounts for nearly 60–70% of the total cost of milk production, making it the largest expenditure in dairy farming. In recent years, the prices of concentrate feeds, oilseed cakes, and fodder have increased considerably, which has reduced the profit margin of farmers and affected the sustainability of dairy enterprises.

Under these conditions, adopting scientific and economical feeding strategies has become essential. Proper feeding does not necessarily mean expensive feeding; rather, it involves providing balanced nutrition using locally available and low-cost feed resources. The use of crop residues, agro-industrial by-products, and green fodder grown on the farm can significantly reduce feeding costs. Scientific feeding also improves nutrient utilization, animal health, and milk production efficiency.

Therefore, adopting low-cost and scientific feeding practices is a key step toward improving productivity, reducing production costs, and increasing the overall profitability and sustainability of dairy farming in India.

Importance of Balanced Ration in Dairy Animals

A balanced ration is essential for maintaining the health, productivity, and efficiency of dairy animals. It provides all the necessary nutrients, including energy, protein, minerals, and vitamins, in the correct proportion required for maintenance, growth, reproduction, and milk production. Energy is needed for daily body functions and milk synthesis, while protein supports muscle development, milk protein formation, and overall body repair. Minerals and vitamins play a vital role in bone development, metabolic activities, immunity, and reproductive performance.

Feeding a balanced ration helps prevent both overfeeding and underfeeding. Overfeeding, especially concentrates, increases feed costs and may lead to metabolic disorders such as acidosis, while underfeeding results in poor growth, low milk yield, and weak animals. Therefore, providing the right quantity and quality of nutrients is important for optimum performance.

Balanced feeding also improves feed efficiency, which means animals utilize nutrients more effectively for milk production rather than wasting them. This leads to higher milk yield,

better animal health, and reduced feeding cost per unit of milk produced. Thus, balanced ration is a key component of scientific and economical dairy farming.

Use of Locally Available Feed Resources

The use of locally available feed resources is an effective and economical strategy to reduce feeding costs in dairy farming. Many farmers can utilize feed materials that are easily available on their farms or in nearby areas, which reduces dependence on expensive commercial feeds. Proper use of these resources not only lowers feeding expenses but also supports sustainable dairy production.

Crop residues such as wheat straw and paddy straw are commonly available after harvesting cereal crops. Although these residues are low in protein, they can serve as a good source of roughage and help maintain proper rumen function. Their nutritive value can be further improved through proper treatment and supplementation with concentrate feed and mineral mixture.

Agro-industrial by-products such as mustard oil cake, groundnut cake, and rice bran are excellent and economical sources of protein and energy. These feed ingredients help meet the nutritional requirements of dairy animals at a lower cost compared to commercial concentrate mixtures.

Seasonal green fodder such as maize, sorghum, berseem, and cowpea provides essential nutrients, improves digestibility, and enhances milk production. Therefore, efficient utilization of locally available feed resources can significantly reduce feed costs and improve farm profitability.

Fodder Production at Farm Level

Fodder production at the farm level is one of the most effective ways to reduce feeding costs and ensure a regular supply of nutritious feed for dairy animals. Farmers can grow green fodder crops such as berseem, maize, sorghum, cowpea, and napier grass depending on the season and local climatic conditions. These fodder crops are rich in essential nutrients and help support better growth, milk production, and overall animal health. Green fodder is also more economical compared to purchased feed and improves feed intake and digestibility.

In addition to green fodder production, farmers should practice fodder conservation methods such as silage and hay making. Silage is prepared from green fodder like maize and sorghum and can be stored for several months without losing much of its nutritional value. Hay is made by drying fodder crops and is useful during periods of green fodder scarcity.

Producing and conserving fodder at the farm level reduces dependency on costly market feeds and ensures year-round availability of quality fodder. This helps lower feeding costs and improves the profitability and sustainability of dairy farming.

Use of Total Mixed Ration (TMR) in Dairy Animals

Total Mixed Ration (TMR) is a scientific feeding method in which roughage (green fodder, dry fodder) and concentrate feed are mixed thoroughly in the proper proportion and offered to dairy animals as a single complete feed. This ensures that animals receive all the essential nutrients, including energy, protein, fiber, minerals, and vitamins, in every bite. TMR can be prepared using locally available feed resources such as green fodder, straw, oilseed cakes, bran, and mineral mixture, making it both economical and efficient.

One of the major advantages of TMR is that it prevents selective feeding by animals. Normally, animals tend to eat more concentrate and leave roughage, which leads to an

imbalance of nutrients and digestive problems. However, in TMR, all feed ingredients are uniformly mixed, so animals cannot separate them, ensuring balanced nutrient intake.

TMR also improves nutrient utilization and digestion, which enhances feed efficiency. As a result, animals produce more milk with the same amount of feed. In addition, TMR reduces feed wastage, improves animal health, and increases overall productivity, making it an effective low-cost feeding strategy for dairy farmers.

Use of Agro-Industrial By-Products in Dairy Animal Feeding

Agro-industrial by-products are valuable and economical feed resources for dairy animals. These by-products are obtained during the processing of crops and agricultural products and can be effectively used as a source of nutrients. Common examples include mustard oil cake, rice bran, wheat bran, and molasses, which are easily available in many parts of India at a lower cost compared to commercial concentrate feeds. These feed ingredients are rich in protein, energy, and essential nutrients required for milk production and animal growth. Mustard oil cake is an excellent source of protein and helps in improving milk yield and body condition. Rice bran and wheat bran provide energy, fiber, and minerals, and also improve digestion. Molasses is a good source of energy and enhances the palatability of feed, encouraging animals to consume more.

The use of these by-products in animal feeding significantly reduces the dependence on expensive commercial concentrates, thereby lowering the overall feeding cost. At the same time, they help maintain proper nutrition, improve feed efficiency, and support better milk production. Therefore, agro-industrial by-products play an important role in economical and sustainable dairy farming.

Use of Urea Treatment of Straw in Dairy Feeding

In India, crop residues such as wheat straw and paddy straw are commonly used as roughage for dairy animals. However, these straws are low in protein and have poor digestibility, which limits their nutritional value and reduces animal productivity. Urea treatment is a simple and cost-effective method to improve the nutritional quality of straw. In this process, straw is treated with a urea solution and stored for a specific period, which increases its nitrogen content and improves its feeding value.

Urea treatment enhances the crude protein content of straw and breaks down complex fiber components, making it more digestible for rumen microorganisms. This improves the utilization of straw and allows animals to obtain more nutrients from low-quality roughage. As a result, animals maintain better body condition and productivity even when good quality fodder is limited.

One of the major benefits of urea-treated straw is that it reduces the requirement for expensive concentrate feed, thereby lowering feeding costs. It also ensures better utilization of locally available crop residues. Thus, urea treatment of straw is an effective and economical feeding strategy for improving dairy animal productivity and farm profitability.

Use of Silage and Hay in Dairy Animal Feeding

Silage and hay are important methods of fodder conservation that help dairy farmers ensure a continuous supply of nutritious feed throughout the year. During seasons when green fodder is abundant, such as the rainy and winter seasons, excess fodder can be preserved as silage or hay. Silage is prepared by fermenting green fodder like maize, sorghum, or napier grass under anaerobic conditions, while hay is made by drying green fodder to reduce its

moisture content. Both methods help retain the nutritional value of fodder for future use.

Silage Preparation Steps (Simple Method)

Step 1: Select suitable fodder such as maize, sorghum, or napier grass.

Step 2: Harvest the crop at proper stage (milk to dough stage in maize).

Step 3: Chop fodder into small pieces (2–3 cm).

Step 4: Fill the chopped fodder in a pit or silo and press it properly to remove air.

Step 5: Cover the pit with polythene sheet and seal with soil to make it airtight.

Step 6: Allow fermentation for 45–60 days.

Step 7: Silage is ready for feeding. Good silage has a pleasant smell and greenish-yellow color

These preserved fodders are especially useful during periods of green fodder scarcity, such as summer or drought conditions. Feeding silage and hay helps maintain animal health, milk production, and growth even when fresh fodder is not available. In addition, fodder conservation reduces dependency on expensive market feeds and ensures efficient use of available resources. Therefore, silage and hay making are economical and essential practices for sustainable and profitable dairy farming.

Mineral Mixture Supplementation in Dairy Animals

Mineral mixture supplementation is a simple and low-cost practice that provides significant benefits to dairy animals. Minerals such as calcium, phosphorus, magnesium, zinc, copper, and iodine are essential for various body functions, including bone development, milk production, reproduction, and immune response. However, these minerals are often deficient in regular feed and fodder, which can negatively affect animal health and productivity.

Providing a balanced mineral mixture daily helps improve milk yield, reproductive efficiency, and disease resistance. It also supports proper growth and maintains overall health. Mineral supplementation prevents deficiency-related problems such as weak bones, low fertility, reduced milk production, and metabolic disorders.

Although mineral mixture requires only a small investment, it greatly improves feed utilization and animal performance. Regular supplementation ensures better productivity, healthier animals, and higher economic returns. Therefore, the use of mineral mixture is an essential component of scientific and low-cost feeding strategies for profitable dairy farming.

Proper Feeding Management Practices in Dairy Animals

Proper feeding management is essential for improving feed efficiency, reducing feeding costs, and maintaining animal productivity. Dairy animals should be fed according to their body weight, milk yield, and physiological status to ensure they receive adequate nutrients without overfeeding or underfeeding. High-yielding animals require more nutrients compared to low-yielding or dry animals, so feeding should be adjusted accordingly.

Avoiding feed wastage is another important aspect of economical feeding. Feed should be offered in proper amounts and clean feeding troughs to prevent spoilage and loss. Providing clean and fresh drinking water at all times is also essential, as water plays a vital role in digestion, nutrient absorption, and milk production.

Maintaining a regular feeding schedule helps animals develop consistent eating habits and improves digestion and nutrient utilization. Good feeding management not only improves animal health and milk production but also reduces unnecessary feed expenses. Therefore, proper feeding practices are essential for efficient and profitable dairy farming.

Table 1: Feeding Schedule Table for Dairy Animals

Animal Type	Green Fodder (kg/day)	Dry Fodder (kg/day)	Concentrate (kg/day)	Mineral Mixture (g/day)	Water
Calf (3–6 months)	2–3	1–2	0.5–1.0	25–30	Free access
Heifer	5–8	2–3	1–2	30–40	Free access
Lactating cow (8–10 L milk)	15–20	4–5	3–4	50–60	Free access
High-yielding cow (>15 L milk)	20–25	5–6	5–6	60–80	Free access
Dry animal	8–10	4–5	1–2	40–50	Free access

Role of Unconventional Feed Resources in Dairy Animal Feeding

Unconventional feed resources are low-cost and locally available materials that can be used as alternative feed ingredients for dairy animals. These include azolla, tree leaves such as subabul, neem, and babul (in limited quantity), and vegetable wastes from households or markets. These feed resources are rich in nutrients and can partially replace expensive concentrate feeds, helping farmers reduce feeding costs.

Azolla is a highly nutritious aquatic plant rich in protein, minerals, and vitamins, and it can be easily grown on the farm with minimal investment. Tree leaves provide protein, fiber, and essential nutrients, while vegetable wastes are a good source of energy and minerals when fed fresh and clean.

The use of these unconventional feed resources helps reduce dependence on costly commercial feeds and improves farm profitability. It also promotes efficient utilization of locally available materials. Therefore, incorporating unconventional feeds in limited and proper amounts is an effective strategy for economical and sustainable dairy farming.

Economic Benefits of Low-Cost Feeding in Dairy Farming

Adopting low-cost feeding strategies provides significant economic benefits to dairy farmers by reducing the overall cost of milk production. Feed is the largest expense in dairy farming, so using locally available feed resources, agro-industrial by-products, and conserved fodder helps lower feeding costs without compromising animal nutrition. This improves the efficiency of resource utilization and reduces dependence on expensive commercial feeds.

Proper and balanced low-cost feeding also improves milk production by ensuring that animals receive adequate nutrients for maintenance and milk synthesis. Healthy animals utilize feed more efficiently, which leads to better growth, improved reproductive performance, and fewer disease problems. This reduces veterinary expenses and improves overall herd performance.

As a result, farmers can produce more milk at a lower cost, increasing their profit margin. Low-cost feeding strategies not only improve farm income but also make dairy farming more sustainable and economically viable. Therefore, scientific and economical feeding practices are essential for enhancing productivity and profitability in dairy farming.

Conclusion

Feed is the major component of dairy farming expenses, but it can be effectively reduced by adopting scientific and economical feeding practices. The use of balanced ration, locally available feed resources, agro-industrial by-products, and conserved fodder such as silage and hay helps lower feeding costs without affecting animal productivity. Proper feeding management, mineral supplementation, and the use of unconventional feed resources further improve nutrient utilization and animal performance.

Careful planning and efficient use of available resources not only reduce feed expenses but also improve milk production, animal health, and reproductive efficiency. This ultimately increases farm income and ensures better economic returns for dairy farmers.

Therefore, low-cost feeding strategies play a crucial role in improving the profitability and sustainability of dairy farming in India. By adopting these practices, farmers can enhance productivity, reduce production costs, and achieve long-term success in dairy farming.